

Certificate of Analysis

Cell count	2 million per vial
Viability	85%
CD8+	94.4%
CD8+NegativeTetramer+	2.0%
CD8+CMVpp65Tetramer+	57.8%
Sterility	Negative for Bacteria, Yeast and Fungi



Donor Information

Donor ID	153
Age	32
Gender	Female
Race	Caucasian
Height	5'7"
Weight	205
ABO Type	O Positive

HLA typing


	Allele 1	Allele 2
HLA-A	*0201	*3101
HLA-B	*15	*57
HLA-C	*03	*06
HLA-DRB1	*1501	*1301

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.



Neg tetramer PE = HLA-A*0201-PE iTag MHC Tetramer (MBL International, Woburn, MA); CMV A2 tetramer = CMV pp65 peptide (NLVPMVATV)/ HLA-A*0201-PE iTag MHC Tetramer (MBL International, Woburn, MA)

IFN-gamma Secretion



20,000 T cells were plated post-thaw in a 96-well round-bottom plate in the presence of 20,000 T2 cells, a B-LCL expressing HLA-A*0201) alone or in the presence of increasing concentrations OF HLA-A*0201 restricted CMV pp65 peptide (NLVPMVATV) or HLA-A*0201 restricted MART-126-35 (A27L) peptide (ELAGIGILTV). Culture media used in this assay is RPMI 1640 + 10% FBS. After an overnight (18-24h) incubation period, supernatant was collected from each well. IFN-γ concentration was analyzed using the Meso Scale Discovery IFN-γ assay.

FOR RESEARCH USE ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.